

Are all gamma-ray-bursts like GRB 980425, GRB 030329 and GRB 031203?

Dafne Guetta¹, Rosalba Perna², Luigi Stella³ and Mario Vietri⁴

ABSTRACT

We study the probability that three GRBs (980425, 030329, 031203) are found within $z = 0.17$, given the luminosity functions consistent with the $\log N - \log S$ relationship for classical cosmological bursts (*i.e.*, those observed by BATSE). We show that, in order for the probability of these three low- z events to be non-negligible (thus making it more likely that they belong to the same class of the classical cosmological bursts), the bursts' luminosity function must be a broken powerlaw. By reasoning in analogy with beamed AGNs, we show that observations are consistent with the expectations if GRB 980425 and GRB 031203 are indeed normal bursts seen sideways. Within this model, no bright burst within $z = 0.17$ should be observed by a HETE-like instrument within the next ~ 20 yr.

Subject headings: gamma rays: bursts

1. Introduction

After the discovery of the association of SN 2003dh with GRB 030329 (Stanek *et al.*, 2003; Hjorth *et al.*, 2003), and of SN 1998bw with 980425, it has been widely accepted that classical, cosmological GRBs arise from the simultaneous collapse of a SN to form a collapsar (Woosley 1994). Further evidence comes from the third nearby burst, GRB 031203, which also has been seen to be closely associated with a type Ic SN (Tagliaferri *et al.*, 2004; Malesani *et al.*, 2004). This model appears plausible for all long GRBs, even the distant ones, in the light of the similarity between these two objects and GRB 011121/SN2001ke (Bloom *et al.*,

¹Hebrew University of Jerusalem & Weizmann Institute, Israel.

²Department of Astrophysical and Planetary Sciences, University of Colorado at Boulder, Boulder, USA.

³INAF-Osservatorio Astronomico di Roma, Monteporzio Catone (Roma), Italy.

⁴Scuola Normale Superiore, Pisa, Italy

2002; Garnavich et al. 2003), an object located at $z \simeq 0.36$. Still, the fact that only some bursts’ afterglows display the re-bumps now associated with the emergence of the optical contribution due to the underlying SN beckons the question of exactly which fraction of all GRBs (within the detectability range of SN rebumps, say $z \lesssim 1$) are in fact associated with simultaneous SNe. In order to tackle this problem, we study whether the detection of GRB 980425, 030329 and 031203, the only ones unequivocally associated with a Type Ic SN of extreme properties within a very small distance ($z=0.17$) from us, is a statistical anomaly, for the luminosity functions that are consistent with the GRB logN-logS distribution.

2. Probabilities of detecting low- z events

If the three low-redshift GRBs (980425, 030329 and 031203) are really typical of the global GRB population, then their discovery within the current time and sky coverage, must be consistent with the local GRB explosion rate as deduced from the very large GRB samples made available by BATSE. A simple computation of the expected number of events within $z = 0.17$ (the redshift of GRB 030329) shows that this is not the case. The volume out to $z = 0.17$, in a $\Omega_\Lambda = 0.7, \Omega = 1$ cosmology with $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$, is $V = 1.4 \text{ Gpc}^3$. Association of a GRB with a SN requires an accurate position, and thus only bursts revealed by the BeppoSAX WFC cameras, or HETE2 WXM telescopes, can be used. Inclusion of INTEGRAL would only make matters slightly worse. These two X-ray telescopes monitor $S_B = 0.123$ and $S_H = 0.806$ steradians respectively (Band, 2002). The total effective operation times for BeppoSAX and HETE2 are generously estimated as $T_B = 4 \text{ yr}$ and $T_H = 2 \text{ yr}$ (L. Piro, private communication). If we call $\dot{n}_o$ the observed local rate of GRBs, we find that the total expected number of GRBs located inside $z = 0.17$ is

$$N = \dot{n}_o V \frac{S_B T_B + S_H T_H}{4\pi} = 0.12 \frac{\dot{n}_o}{0.5 \text{ Gpc}^{-3} \text{ yr}^{-1}} \quad (1)$$

The probability of observing three bursts, assuming Poisson statistics, is then $P_3 = 2.7 \times 10^{-4} (\dot{n}_o / (0.5 \text{ Gpc}^{-3} \text{ yr}^{-1}))^3$.

The major uncertainty in the above equation concerns $\dot{n}_o$, which takes on different values depending upon exactly which data property is fitted (V/V_{max} , or the $\log N - \log P$ relationship for BATSE data). For the local observed rate, Porciani and Madau (2001) consider three different star formation rates, and correspondingly find three values for $\dot{n}_o$, between 0.11 and 0.17 bursts $\text{Gpc}^{-3} \text{ yr}^{-1}$ (having converted their results to $H_0 = 71 \text{ km s}^{-1} \text{ Mpc}^{-1}$). Schmidt (2001) considers the same star formation rates, but fits different quantities, to obtain $\dot{n}_o = 0.48, 0.51, 0.72 \text{ Gpc}^{-3} \text{ yr}^{-1}$. Perna, Frail and Sari (2003) find $\dot{n}_o = 0.5 \text{ Gpc}^{-3} \text{ yr}^{-1}$, while Guetta, Piran and Waxman (2003) find $\dot{n}_o = 0.44 \text{ Gpc}^{-3} \text{ yr}^{-1}$. In the next

section, we will repeat the analysis of Guetta *et al.*(2003) with a more updated sample, to obtain $\dot{n}_o = 1.1 \text{ Gpc}^{-3} \text{ yr}^{-1}$. With this value, the probability of a triple event would be $P_3 = 2.9 \times 10^{-3}(\dot{n}_o/(1.1 \text{ Gpc}^{-3} \text{ yr}^{-1}))^3$. The hypothesis that the three nearby bursts, 980425, 030329 and 031203, belong to the same group as the classical BATSE bursts, would thus be rejected with fairly high confidence.

However, this is not correct. The local rate discussed above has been derived by all authors under the hypothesis that classical bursts exceed by far the luminosity of GRB 980425 and they also exceed that of GRB 031203 (i.e. they considered a minimum luminosity for the LF $L_b \gtrsim 5 \times 10^{49} \text{ erg/sec}$, which from now on we will assume to define the minimum luminosity for the classical population). Therefore, the derived probabilities are not self-consistent. An unified picture can only be obtained with a LF that includes all luminosities down to that of GRB 980425, and at the same time yields high probability of observing the three low- z events. This is what we will achieve in §4, after setting up the formalism and recomputing (for the more updated sample) the local rate for the classical population in §3.

3. Luminosity function for classical GRBs

We consider all 2204 GRBs of the GUSBAD catalog which contains all the long GRBs ($T_{90} > 2\text{sec}$) (Kouveliotou et al., 1993), detected while the BATSE onboard trigger (Paciesas et al. 1999) was set for 5.5σ over background (resolution of 1024 ms) in at least two detectors, in the energy range 50-300 keV. We estimate $C_{\text{max}}/C_{\text{min}}$ for each burst (where C_{max} is the count rate in the second brightest illuminated detector and C_{min} is the minimum detectable rate) and find $\langle V/V_{\text{max}} \rangle = 0.335 \pm 0.007$.

The luminosity function (LF) of the “classical” population of long-duration GRBs may be represented as a power law with lower and upper limits, L_b and L_2 , respectively. The local LF of GRB peak luminosities L , defined as the co-moving space density of GRBs in the interval $\log L$ to $\log L + d \log L$ is:

$$\Phi_o(L) = c_o(L/L_b)^{-\alpha}, \quad L_b < L < L_2 \quad (2)$$

where c_o is a normalization constant. We stress that this LF is the “isotropic-equivalent” LF, it does not include the effects of beaming.

We assume that GRBs trace the star-formation history, and adopt the recent SFR derived by Rowan-Robinson (1999; RR-SFR); this can be fitted with the expression $R_{\text{GRB}}(z) = \dot{n}_o \max(0.75, 0.75 z)$

The modeling procedure involves the derivation of the peak flux $P(L,z)$ of a GRB of

peak luminosity L observed at redshift z :

$$P(L, z) = \frac{L}{4\pi D_L^2(z)} \frac{C(E_1(1+z), E_2(1+z))}{C(E_1, E_2)}, \quad (3)$$

where $D_L(z)$ is the bolometric luminosity distance and $C(E_1, E_2)$ is the spectral energy distribution integrated between $E_1 = 50$ keV and $E_2 = 300$ keV. Schmidt (2001) finds that the median value of the spectral photon index in the 50-300keV band for long bursts GRBs is -1.6. We use this value for our analysis to include the k-correction.

Objects with luminosity L observed by BATSE with a flux limit P_{lim} are detectable to a maximum redshift $z_{\text{max}}(L, P_{\text{lim}})$ that can be derived from Eq. (3). We consider an average limiting flux $P_{\text{lim}} = 0.25 \text{ ph cm}^{-2}\text{s}^{-1}$ taken from the BATSE catalog.

The number of bursts with a peak flux $> P$ is given by:

$$N(> P) = \int \Phi_o(L) d \log L \int_0^{z_{\text{max}}(L, P)} \frac{R_{\text{GRB}}(z)}{1+z} \frac{dV(z)}{dz} dz \quad (4)$$

where the factor $(1+z)^{-1}$ accounts for cosmological time dilation and $dV(z)/dz$ is the comoving volume element.

To be consistent with previous calculations, we take the low luminosity cut-off $L_b \sim 5 \times 10^{49} \text{ erg s}^{-1}$, while the high luminosity, $L_2 \sim 5 \times 10^{52} \text{ erg s}^{-1}$, is taken on the order of the maximum luminosity detected until the time of writing (Bloom et al. 2003). The slope α of the LF is constrained by fitting the model with the observed peak flux distribution with a non-linear Levenberg-Marquart minimum χ^2 method. To avoid problems with error correlation in a cumulative distribution like $N(> P)$ propagate in an unknown way, we use the differential distributions $n(P) \equiv dN/dP$ for this analysis. We find that the best fit is given for $\alpha \sim 0.72$; the p-value of the fit is ~ 0.70 (see Fig.1 upper panel). Both α and the normalization value are somewhat insensitive to the choice of L_b below a value $\sim 10^{50} \text{ erg s}^{-1}$. This is mainly because GRBs with very low luminosity appear above the sensitivity limit of $\sim 0.25 \text{ photons cm}^{-2} \text{ s}^{-1}$ in a very small volume around the observer.

To obtain the observed local rate of GRBs per unit volume, $\dot{n}_o$, we need to estimate the effective full-sky coverage of our GRB sample. The GUSBAD catalog represents 3.185 years of BATSE full sky coverage implying a rate of 692 GRB per year. Using our LF we find $\dot{n}_o \sim 1.1 \text{ Gpc}^{-3}\text{yr}^{-1}$.¹

¹Note that for a different SFR like for example the SF2-SFR of Porciani and Madau (2001) we still find

4. Luminosity function for bursts seen from any direction and revised probabilities

We are thus left with the question of how to include GRB 980425 and GRB 031203 within our analysis. It is quite possible that these bursts belong to a different, local population of bursts, as occasionally claimed by some authors for GRB 980425 (Bloom et al. 1998). Still, there are at least two arguments suggesting that 980425 is just a normal burst seen sideways (Nakamura 1998; Eichler & Levinson 1999; Woosley, Eastman & Schmidt 1999). One is the exceptional similarity of the two underlying SNe in 980425 and 030329, within a class (that of extreme Type Ic SNe) known instead for its lack of common patterns. The other is that GRB 021211 (della Valle *et al.*, 2003), and all bursts with the so-called optical rebumps are likely associated also to SN.

For this reason, we consider a LF model which includes GRBs seen sideways. We follow the analogy with beamed AGNs, according to which (Urry and Shafer 1984) the total LF is simply a broken power-law, agreeing, at the bright end, with the LF of the objects seen face-on. We thus take:

$$\Phi_o(L) = c_o \begin{cases} (L/L_b)^{-\gamma}, & L_1 < L < L_b \\ (L/L_b)^{-\alpha}, & L_b < L < L_2 \end{cases} \quad (5)$$

We impose here $L_1 = L_{980425}$, $L_2 = 5 \times 10^{52}$ erg s⁻¹ and L_b to be the border value between GRBs seen sideways (weak bursts) and the ones seen face-on (classical bursts). We choose somewhat arbitrarily $L_b \sim 5 \times 10^{49}$ erg s⁻¹, and fit the same data as in the previous section with α and γ as free parameters. Note that it has been proposed that X-ray flashes (XRFs) are simply GRBs viewed off-axis. If this interpretation is correct, then XRFs could be included in our analysis in the lower part of the LF broken power law (recalling that XRFs are generally underluminous compared to GRBs). The best-fit ($\alpha = 0.7$, $\gamma = 0.1$ and p-value ~ 0.76) is shown in Fig. 1 (lower panel).² With this LF, the total local rate of events, down to the lowest luminosity (that of 980425) is now much higher, $\dot{n}_o = 10$ Gpc⁻³yr⁻¹, while the rate for $L > L_b$ remains unaffected. Using Eq(1), we find that the expected number of events within $z = 0.17$, and of luminosity even as low as GRB 980425, is $N_l = 2.4$, which is clearly highly compatible with the current observation of 3 events.

a good fit and a local rate smaller by a factor ~ 2 . We find a similar result also for the LF studied in the next session.

²Note that the curve is almost identical to the one corresponding to the best fit LF obtained for the classical bursts shown in the upper panel. This is because to add a low luminosity ($L < 5 \times 10^{49}$ erg/sec) tail to the LF does not affect the logN-logS distribution (low luminosity GRBs appear above the BATSE sensitivity limit in a reduced volume as explained in section 4).

A strong radio emission is expected from bursts seen sideways at ~ 1 yr delay (Waxman 2004ab, Livio & Waxman 2000, see also Waxman 2004a,b and reference within) as the jet decelerates to sub-relativistic speed and its emission approaches isotropy. However no late-time rebrightening was detected for either GRB 980425 or GRB 031203 (Sodeberg, Frail & Wieringa 2004; Sodeberg et al. 2003, 2004). These results suggest that neither GRB 980425 nor GRB 031203 were off-axis events, and instead were intrinsically sub-energetic GRBs. Therefore we also consider the possibility that both GRB 980425 and GRB 031203 might have intrinsically low luminosity and all GRBs are seen on axis. The LF can still have an *intrinsic* break for some (unknown) reason, and therefore it can still be represented by eq.(5) (as in Schmidt 2001). In this case we can leave L_b as a free parameter in our fit and look for the best fit parameters α, γ, L_b . The results ($\alpha = 0.95$, $\gamma = 0.4$, $L_b = 10^{51}$ erg s^{-1} and p-value ~ 0.64) are shown in Fig. 1 lower panel. With this LF the total local rate of events is $\dot{n}_o = 20$ Gpc $^{-3}$ yr $^{-1}$. However, if all GRBs are seen on axis, then there is no natural explanation for the break in the LF. Indeed a single power law LF with $L_1 = L_{980425}$, $L_2 = 5 \times 10^{52}$ erg s^{-1} could be representative of these bursts. However in this case we get a very high local rate, $\dot{n}_o \sim 100$ Gpc $^{-3}$ yr $^{-1}$, which would predict about 24 events within the $z = 0.17$ volume (again using Eq.1). The probability of having only 3 occurrences is $\sim 10^{-7}$, which makes this scenario very unlikely. Therefore, we conclude that a broken powerlaw for the luminosity function is highly preferred to a single powerlaw, if the three low- z events do belong to the same class as the classical ones. Also note that the rate for the single powerlaw would yield about 400 GEM (galactic events per million years), implying an implausibly large number of GRB remnants (Loeb & Perna 1998; Efremov et al. 1998) at any given time.

Our model also allows us to compute the probability of finding one object (GRB 980425) as close as $z = 0.008$, which would be of course prohibitively small if we were to apply to this burst the statistics for classical GRBs discussed in §3. For our best-fit powerlaw model, we find that the probability of bursts as luminous as GRB 980425 within the volume accessible to BeppoSAX and HETEII, is $\approx 10^{-3}$, small but still not completely ruled out. For the detection threshold of Swift (Gehrels et al. 2004), this rate would be 0.05, while that of all bursts with $L < L_b$ is ~ 2 for a 2-yr observation period.




Fig. 1.— Upper panel: The best-fit differential distribution of the classical GRBs vs. the observed one taken from the GUSBAD catalog. Lower panel: The best-fit differential distribution of the GRBs in the case in which L_b marks the border between off-axis and on axis-GRBs (solid line) and in the case in which we leave L_b as a free parameter (dashed line) vs. the observed one taken from the GUSBAD catalog. For both panels: The $\log_{10}(C_M/C_m)$ is divided into 14 bins, the first 11 of equal size and the remaining 3 with varying sizes chosen such that the number of bursts per bin in the observed sample is at least $N_{\min} = 40$ (in order to have reasonable statistics, so that the Poisson error will not be too large), and N is the observed (*circles*) or theoretical (*lines*; only the values at the center of each bin count) number of bursts in each bin. The edges of the bins are also plotted.

5. Conclusions

We have shown in this *Letter* that the presence of three nearby GRBs does not pose a problem for the current view (according to which both distant and nearby bursts belong to the same class), if we modify the LF to include an extension to luminosities as low as that of GRB 980425. We did this by means of a *minimum impact* extension, *i.e.*, by assuming, in complete analogy with beamed AGNs, that the LF of the bursts is a broken power-law, with the bright-end distribution equal to what we derived when we neglected the presence of faint, sideways bursts. As a by-product, we showed that this yields a non-negligible probability for the detection of GRB 980425-like event, which is also a new result.

How could this picture change? Some authors have insisted that it is not possible to throw GRB 980425 in the same cage as classical GRBs, because of its unique properties in the radio band. Within our model, the local rate of nearby bright bursts ($L \gtrsim 5 \times 10^{49}$ erg s $^{-1}$) observable by HETEII within $z = 0.17$ is $\dot{n}_o V S_H / 4\pi = 0.057$ yr $^{-1}$. This implies that we ought to observe the next such event ~ 20 yr from now. Should we see one significantly earlier than that then the argument suggesting a similarity between the distant, classical bursts and the nearby ones would have to be reassessed.

We thank the anonymous referee for his/her very useful comments. We thank Davide Lazzati for his comments. DG acknowledges the NSF grant AST-0307502 for financial support.

REFERENCES

- Band, L. 2002, ApJ 572, L45.
- Bloom, J. S., Kulkarni, S. R., Harrison, F., Prince, T., Phinney, E. S., Frail, D. A. 1998, ApJ, 506L, 105
- Bloom, J. S., Frail, D. A., Kulkarni, S. R. 2003, ApJ 594, 674.
- Kouveliotou, C. et al. 1993, ApJ 431, 101.
- Della Valle, M. et al. 2003, A&A 406, L33.
- Eichler, D., & Levinson, A. 1999, ApJ 521, L11.
- Efremov, Y. N., Elmegreen, B. G., & Hodge, P. W. 1998, ApJ, 501, L163
- Garnavich P. M. et al., 2003, ApJ 582, 924.
- Gehrels, G. et al., 2004, ApJ 611, 1005.
- Guetta, D., Piran, T. and Waxman, E. astro-ph/0311488, ApJ in press.

- Hjorth et al., 2003, *Nature* 423, 847.
- Livio, M. & Waxman, E. 2000, *ApJ* 538, 187.
- Loeb, A., & Perna, R. 1998, *ApJ*, 533, L35
- Malesani, D. et al. 2004, astro-ph/0405449 *ApJ Letters* in press.
- Nakamura, T. 1998, *Prog. Theor. Phys.*, 100, 921.
- Paciesas, W. et al. 1999, *ApJS* 122, 456.
- Perna, R., Sari, R., Frail, D. 2003, *ApJ* 594, 379.
- Porciani, C. & Madau, P. 2001, *ApJ* 548, 522.
- Schmidt, M. 2001, *ApJ* 552, 365.
- Soderberg, A. M., Frail, D. A. & Wieringa, M. H., 2004, *ApJL* 607, L13.
- Soderberg, A. M. et al., 2004, *Nature*, in press.
- Soderberg A. M. et al., 2003, *GCN* 2483.
- Stanek et al. 2003, *ApJ* 591, L17.
- Tagliaferri, G. et al. 2004, *IAU Circ.*, 8308, 1. Edited by Green, D. W. E.
- Urry, C. M. & Shafer, R. A. 1984, *ApJ* 280, 569.
- Waxman, E. 2004a, *ApJ* 605, L97.
- Waxman, E. 2004b, *ApJ* 602, 886.
- Woosley, S. E. 1994, *Bulletin of the American Astronomical Society*, Vol. 26, p.1413.
- Woosley, S. E., Eastman, R. & Schmidt, B. P. 1999, *ApJ*, 516, 788.